

Roll No.

Total No. of Pages : 02

Total No. of Questions : 09

B.Tech. (3D Animation & Graphics) (2012 Onwards)

B.Tech. (CSE)/(IT) (2012 Batch)

(Sem.-3)

MATHEMATICS-III

Subject Code : BTAM-302

Paper ID : [A2143]

Time : 3 Hrs.

Max. Marks : 60

INSTRUCTION TO CANDIDATES :

1. SECTION-A is COMPULSORY consisting of TEN questions carrying TWO marks each.
2. SECTION-B contains FIVE questions carrying FIVE marks each and students has to attempt any FOUR questions.
3. SECTION-C contains THREE questions carrying TEN marks each and students has to attempt any TWO questions.

SECTION-A

1. Write briefly :

- (a) Define periodic function. Give an example of a function which is not periodic.
- (b) Write the sufficient conditions for the existence of Laplace transform.
- (c) Find the Laplace transform of $\cos(at)$.
- (d) Obtain a partial differential equation that governs the family of surfaces $z = (x - \alpha)^2 + (y - \beta)^2$.
- (e) Define linear partial differential equation and give an example of a partial differential equation which is not linear.
- (f) Write the sufficient conditions for a function of complex variable to be analytic.
- (g) Gauss elimination method is used to solve which equations?
- (h) Write the fourth-order Runge-Kutta method to solve initial value problems of ordinary differential equations.
- (i) The number of emergency admissions each day to a hospital is found to have Poisson distribution with mean 4. Find the probability that on a particular day there will be no emergency admissions.
- (j) Write one application of F-distribution.

SECTION-B

2. Find the Fourier cosine series of the function

$$f(x) = \begin{cases} x^2, & 0 \leq x \leq 2, \\ 4, & 2 \leq x \leq 4. \end{cases}$$

3. State and prove linearity property of Laplace transform and use it to find the Laplace transform of $\sinh(3t)$ and $\cosh(4t)$.
4. Solve : $(x^2 - yz)p + (y^2 - zx)q = z^2 - xy$, where $p = \frac{\partial z}{\partial x}$; $q = \frac{\partial z}{\partial y}$.
5. Two players A and B play tennis games. Their chances of winning a game are in the ratio 3 : 2 respectively. Find A's chance of winning at least two games out of four games played.
6. Two random samples of sizes 9 and 7 gave the sum of squares of deviations from their respective means as 175 and 95 respectively. Can they be regarded as drawn from normal populations with the same variance ?

SECTION-C

7. If $f(z) = u + iv$ is an analytic function of $z = x + iy$ and $u + v = (x + y)(2 - 4xy + x^2 + y^2)$, then find u , v and the corresponding analytic function $f(z)$.
8. Find the solution of the system of equations

$$45x_1 + 2x_2 + 3x_3 = 58$$

$$-3x_1 + 22x_2 + 2x_3 = 47$$

$$5x_1 + x_2 + 20x_3 = 67$$

correct to three decimal places using the Gauss-Seidel iteration method.

9. Solve the initial value problem $\frac{dy}{dx} = -2xy^2$; $y(0) = 1$ with the step size $h = 0.2$

on the interval $[0, 0.6]$ using the classical fourth-order Runge-Kutta method. The

exact solution of the problem is $y(x) = \frac{1}{1+x^2}$. Find the absolute errors at each

step.